

A NOTE ON SIMPLICIAL DIMENSION SHIFTING

ADRIAN OCNEANU

ABSTRACT. We discuss a simplicial dimension shift which associates to each n -manifold an $n-1$ -manifold. As a corollary we show that an invariant which was recently proposed by Ooguri and by Crane and Yetter for the construction of 4-dimensional quantum field theories out of 3-dimensional theories is trivial.

Ooguri [1] proposed recently a method for obtaining 4-manifold invariants and studying quantum gravity, with a construction inspired statistical mechanics. This procedure was further formalized by Crane and Yetter [2]. Their method is associating a 3-manifold with boundary to every 4-simplex Δ^4 . Given 4-manifold M with a triangulation τ and an invariant for 3-manifolds with boundary, they evaluate the invariant on the 3-manifolds with boundary coming from each 4-simplex, they match boundary conditions according to the triangulation and then they compute a partition function. They mention that the proof of the invariance of the result to the choice of the triangulation is long and technical.

In fact the matching and summation of boundary conditions is equivalent to the glueing together of the corresponding portions of the boundary and then computing the invariant of the 3-manifold without boundary obtained this way. With this observation the choice of the 3-manifold invariant becomes irrelevant. One obtains a topological construction, which associates to each triangulated 4-manifold a 3-manifold, which we call the dimension shift of M . The aim of this note is to prove that the dimension shift of a 4-manifold is essentially trivial, and that for a singular triangulated 4-manifold the dimension shift yields essentially the direct sum of the 3-singularities of M . In particular, the invariants of Ooguri, Crane and Yetter are always 1.

We were led independently to the same construction in a different context [3]. We showed an equivalence between systems of n -modules over operator algebras and simplicial invariants for $n+1$ -manifolds which have the glueing property. Working with subfactors, or 2-modules, an asymptotic construction produces naturally some 3-modules. The 4-manifold invariants associated to these 3-modules are

Date: January 1993.

obtained from the 3-manifold invariants associated to the 2-modules by means of dimension shifting. Because of the result which follows we used this construction only to show that the axioms for 4-dimensional invariants are nonempty.

The idea of dimension shifting is the following. Let Δ^4 denote the 4-simplex. The boundary $\partial\Delta^4$ of Δ^4 consists of 5 copies of the 3-simplex Δ^3 , and is topologically equivalent to the 3-sphere S^3 . Consider the 1-skeleton σ_1 of $\partial\Delta^4$ and the dual 1-skeleton $\widehat{\sigma}_1$ of $\partial\Delta^4$. Let T be a small tubular neighborhood of $\widehat{\sigma}_1$ in S^3 .

Let now M^4 be a 4-manifold with a triangulation τ . To each 4-simplex $\Delta^4 \in \tau$ associate the 3-manifold with boundary T as above. When two 3-faces Δ^3 and Δ'^3 of the 4-simplices Δ^4 and Δ'^4 in τ are glued, glue the corresponding components $\partial T \cap \Delta^3$ and $\partial T' \cap \Delta'^3$ of the boundary of T and respectively T' .

This way one obtains a 3-manifold without boundary denoted by $\text{sh}_\tau^{\text{int}} M$. In another version of this construction one glues the exterior $S^3 - T$ of T the same way as above, to obtain a 3-manifold $\text{sh}_\tau^{\text{ext}} M$. Note that $S^3 - T$ is a tubular neighborhood of the 1-skeleton σ_1 .

Theorem 0.1. *Let M^4 be a connected compact 4-manifold without boundary, with a triangulation τ having n_k simplices in each dimension $k = 0, \dots, 4$. Then we have*

$$\text{sh}_\tau^{\text{int}} M = (10n_4 - n_3 + 1)(S^1 \times S^2)$$

and

$$\text{sh}_\tau^{\text{ext}} M = (n_1 - n_0 + 1)(S^1 \times S^2).$$

Here mM denotes the connected sum of m copies of the manifold M .

Proof. The proof for the case of the interior dimension shift is the following. Let Δ^4 be a 4-simplex in τ , let T be its thickened dual 1-skeleton and let Δ^3 be a 3-simplex in $\partial\Delta^4$. Then $T \cap \Delta^3$ is topologically a 3-ball, and its boundary consists of $\partial T \cap \Delta^3$ and of $T \cap \partial\Delta^3$, which is the union of 4 copies of a 2-disk D^2 . When Δ^4 is glued to another 4-simplex Δ'^4 in τ , and Δ^3 is identified to its pair Δ'^3 in $\partial\Delta'^4$, the balls $T \cap \Delta^3$ and $T' \cap \Delta'^3$ are glued together to yield a copy of S^3 which has a 3-ball hole B^3 carved out for each face Δ^2 of Δ^3 . The boundary S^2 of the hole B^3 consists of a pair of disks D^2 and D'^2 as described above. The disk D^2 is then glued to a disk coming from the neighbor Δ'^3 of Δ^2 in $\partial\Delta^4$ opposite to Δ^3 .

This procedure yields a 3-sphere for every 3-simplex of τ and a handle $S^2 \times I$ for every 2-face Δ^2 of each 4-simplex Δ^4 of τ , for a total of $10n_4$

handles. From these, $n_3 - 1$ handles are needed to connect the 3-spheres, and $10n_4 - n_3 + 1$ handles remain.

In the case of the exterior shift, note that the exterior $S^3 - T$ corresponding to a 4-simplex Δ^4 is a tubular neighborhood of the 1-skeleton of Δ^4 . We decompose $S^3 - T$ into balls B_v^3 for each vertex v of Δ^4 joined by bars $b_e = D_e^2 \times I$, with I the 1-simplex, obtained by thickening each 1-edge e of Δ^4 .

Let now v be a vertex of the triangulation τ of M^4 . The simplicial neighborhood, or star, of v is a cone over a 3-manifold M_v called the link of v . The link M_v is obtained by joining the faces Δ_i^3 opposed to v in the 4-simplices Δ_i^4 which contain v . For each vertex v , each neighboring 4-simplex Δ_i^4 contributes to $\text{sh}_\tau^{\text{ext}} M$ with a 3-ball B_{vi}^3 .

The crucial observation is that the ball B_{vi}^3 is obtained by a homothety from the 3-simplex Δ_i^3 opposite to v in Δ_i^4 , and the glueing of the 3-balls B_{vi}^3 is carried over by this homothety from the glueing of the corresponding link simplices Δ_i^3 . This yields a 3-manifold $\widetilde{M}_v$ isomorphic to M_v for each vertex v . The manifold $\widetilde{M}_v$ has a hole for each edge e of τ , with boundary obtained by joining the disks D_{vei}^2 which connect the balls B_{vi} to the bars b_{ei} . By the same homotopy, the disks D_{vei}^2 join to make a 2-manifold $\widetilde{M}_e$ which is homothetic to the link M_e of e . The bars themselves glue to give copies of $M_e \times I$.

For a manifold triangulation τ each M_v is a 3-sphere and each M_e is a 2-sphere, and we obtain n_0 spheres S^3 and n_1 handles $S^2 \times I$. A counting argument similar to the one done before ends the proof. $\square$

Remark 0.1. Analogous dimension shifts can be defined from an n -dimensional triangulated manifold M^n to an $n-1$ -manifold, using tubular neighborhoods of the k -skeleton of Δ^n . The proof above extends to this context and shows that the interior dimension shift carries essentially no information at all about M . If M is a triangulated manifold with singularities, i.e. such that the links of M are not all spheres, then the exterior dimension shift describes the link singularities of M . Thus, for the 4-dimensional case described above, the shifted invariant of a singular triangulated 4-manifold is the product of the 3-dimensional invariants of the links of its vertices. This is the result that we described in [3].

REFERENCES

1. H.Ooguri, *Topological lattice methods in four dimensions*, Preprint, Kyoto University.
2. L.Crane and D.Yetter, *A categorical construction of 4-d topological quantum field theories*, Dec. 1992, Preprint, Kansas State University.

3. A. Ocneanu, *Operator algebras, quantum groups and manifold invariants*, Feb. 1992, Colloque, Universite Paris VII.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY,
UNIVERSITY PARK, PENNSYLVANIA 16801, USA